



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

PROBLEMS FOR SOLUTION.

ALGEBRA.

259. Proposed by ARTEMUS MARTIN, M. A., Ph. D., LL. D., Washington, D. C.

On page 167 of George Bruce Halsted's *Metrical Geometry* (Mensuration), Boston, 1881, Table of Scalene Triangles, is found the following triangle, viz., Sides 21, 61, 65; Area 420. The sides of a rational scalene triangle, whose sides have no common divisor, can not all be odd; one must be even and the other two odd. It is required to find the error in the sides of the above triangle, assuming that the area is correct.

260. Proposed by O. E. GLENN, Ph. D., Springfield, Mo.

The necessary and sufficient condition that a binary form be a perfect n th power is that its Hessian vanish.

261. Proposed by REV. R. D. CARMICHAEL, Hartselle, Ala.

Sum to infinity the series, $\frac{1}{n^p} + \frac{3}{n^{2p}} + \frac{5}{n^{3p}} + \frac{7}{n^{4p}} + \frac{9}{n^{5p}} + \dots$

AVERAGE AND PROBABILITY.

176. Proposed by T. N. HAUN, Mohawk, Tenn.

A cube being cut at random by a plane, what is the chance that the section is a hexagon? (Problem 72, p. 503, Williamson's *Integral Calculus*.)

CALCULUS.

217. Proposed by PROFESSOR F. ANDEREGG, Oberlin College, Oberlin, Ohio.

Find $\lim_{n \rightarrow \infty} \frac{1}{n} \sqrt[n]{(n+1)(n+2)\dots(2n)}$.

218. Proposed by REV. R. D. CARMICHAEL, Hartselle, Ala.

Evaluate (a) $\int_0^{\frac{1}{2}\pi} \frac{\sin mx \sin nx}{\sin x} dx$; (b) $\int_0^{\frac{1}{2}\pi} \frac{\cos mx \sin nx}{\sin x} dx$, where n is a positive integer. Also, modify the result for the case of m an integer.

219. Proposed by C. N. SCHMALL, College of the City of New York, New York City.

In the article "Infinitesimal Calculus" in the *Encyclopaedia Britannica* Vol. XIII, page 24, I notice the following: "Of all triangular pyramids standing on a given triangular base, and of given altitude, find that whose surface is the least." A solution is required.